# SUMMARY OF PRODUCT CHARACTERISTICS

### **1** NAME OF THE MEDICINAL PRODUCT

Prednisolone 1 mg Tablets Prednisolone 5 mg Tablets

# 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each Tablet contains 1 mg of Prednisolone. Excipients with known effect: Also contains 36 mg of Lactose.

Each Tablet contains 5 mg of Prednisolone. Excipients with known effect: Also contains 58 mg of Lactose.

For the full list of excipients, see section 6.1.

# **3 PHARMACEUTICAL FORM**

Tablet

Appearance: White, circular, biconvex tablets with "P1" on one side and plain on the reverse.

*Appearance:* White, circular, biconvex tablets with P5 on one side and break line on the reverse The tablet can be divided into equal doses.

# 4 CLINICAL PARTICULARS

#### 4.1 Therapeutic indications

Allergy and anaphylaxis: bronchial asthma, drug hypersensitivity reactions, serum sickness, angioneurotic oedema, anaphylaxis.

Arteritis/collagenosis: giant cell arteritis/polymyalgia rheumatica, mixed connective tissue disease, polyarteritis nodosa, polymyositis.

**Blood disorders:** haemolytic anaemia (auto-immune), leukaemia (acute and chronic lymphocytic), lymphoma, multiple myeloma, idiopathic thrombocytopenic purpura.

**Cardiovascular disorders:** post-myocardial infarction syndrome, rheumatic fever with severe carditis.

**Endocrine disorders:** primary and secondary adrenal insufficiency, congenital adrenal hyperplasia.

**Gastro-intestinal disorders:** Crohn's disease, ulcerative colitis, persistent coeliac syndrome (coeliac disease unresponsive to gluten withdrawal), auto-immune chronic active hepatitis, multisystem disease affecting liver, biliary peritonitis.

Hypercalcaemia: sarcoidosis, vitamin D excess.

**Infections (with appropriate chemotherapy):** helminthic infestations, Herxheimer reaction, infectious mononucleosis, miliary tuberculosis, mumps orchitis (adult), tuberculous meningitis, rickettsial disease.

Muscular disorders: polymyositis, dermatomyositis.

**Neurological disorders:** infantile spasms, Shy-Drager syndrome, sub-acute demyelinating polyneuropathy.

**Ocular disease:** scleritis, posterior uveitis, retinal vasculitis, pseudo-tumours of the orbit, giant cell arteritis, malignant ophthalmic Graves' disease.

**Renal disorders:** lupus nephritis, acute interstitial nephritis, minimal change glomerulonephritis.

**Respiratory disease:** allergic pneumonitis, asthma, occupational asthma, pulmonary aspergillosis, pulmonary fibrosis, pulmonary alveolitis, aspiration of foreign body, aspiration of stomach contents, pulmonary sarcoid, drug induced lung disease, adult respiratory distress syndrome, spasmodic croup.

**Rheumatic disorders:** rheumatoid arthritis, polymyalgia rheumatica, juvenile chronic arthritis, systemic lupus erythematosus, dermatomyositis, mixed connective tissue disease.

Skin disorders: pemphigus vulgaris, bullous pemphigoid, systemic lupus erythematosus, pyoderma gangrenosum.

**Miscellaneous:** sarcoidosis, hyperpyrexia, Behçets disease, immunosuppression in organ transplantation.

#### 4.2 **Posology and method of administration**

#### **Posology**

The initial dosage of Prednisolone may vary from 5 mg to 60 mg daily depending on the disorder being treated. Divided daily dosage is usually used.

The following therapeutic guidelines should be kept in mind for all therapy with corticosteroids:

Corticosteroids are palliative symptomatic treatment by virtue of their antiinflammatory effects; they are never curative.

The appropriate individual dose must be determined by trial and error and must be re-evaluated regularly according to activity of the disease.

As corticosteroid therapy becomes prolonged and as the dose is increased, the incidence of disabling side-effects increases.

In general, initial dosage shall be maintained or adjusted until the anticipated response is observed. The dose should be gradually reduced until the lowest dose which will maintain an adequate clinical response is reached. Use of the lowest effective dose may also minimise side-effects (see Section 4.4 'Special warnings and special precautions for use').

In patients who have received more than physiological dose for systemic corticosteroids (approximately 7.5 mg prednisolone or equivalent) for greater than 3 weeks, withdrawal should not be abrupt. How dose reduction should be carried out depends largely on whether the disease is likely to relapse as the dose of systemic corticosteroids is reduced. Clinical assessment of disease activity may be needed during withdrawal. If the disease is unlikely to relapse on withdrawal of systemic corticosteroids but there is uncertainty about hypothalamic-pituitary-adrenal (HPA) suppression, the dose of corticosteroid may be reduced rapidly to physiological doses. Once a daily dose equivalent to 7.5 mg of prednisolone is reached, dose reduction should be slower to allow the HPA-axis to recover.

Abrupt withdrawal of systemic corticosteroid treatment, which has continued up to 3 weeks, is appropriate if it is considered that the disease is unlikely to relapse. Abrupt withdrawal of doses of up to 40 mg daily of prednisolone or equivalent for 3 weeks is unlikely to lead to clinically relevant HPA-axis suppression, in the majority of patients. In the following patient groups, gradual withdrawal of systemic corticosteroid therapy should be considered even after courses lasting 3 weeks or less:

• Patients who have had repeated courses of systemic corticosteroids, particularly if taken for greater than 3 weeks.

• When a short course has been prescribed within one year of cessation of long-term therapy (months or years).

• Patients who may have reasons for adrenocortical insufficiency other than exogenous corticosteroid therapy.

• Patients receiving doses of systemic corticosteroid greater than 40 mg daily of prednisolone (or equivalent).

• Patients repeatedly taking doses in the evening.

(See Section 4.4 'Special warnings and special precautions for use' and Section 4.8 'Undesirable effects')

During prolonged therapy, dosage may need to be temporarily increased during periods of stress or during exacerbations of the disease (see Section 4.4 'Special warnings and special precautions for use')

If there is lack of a satisfactory clinical response to prednisolone tablets, the drug should be gradually discontinued and the patient transferred to alternative therapy.

**Intermittent dosage regimen** A single dose of prednisolone tablets in the morning on alternate days or at longer intervals is acceptable therapy for some patients. When this regimen is practical, the degree of pituitary-adrenal suppression can be minimised.

**Specific dosage guidelines** The following recommendations for some corticosteroid-responsive disorders are for guidance only. Acute or severe disease may require initial high dose therapy with reduction to the lowest effective maintenance dose as soon as possible. Dosage reductions should not exceed 5-7.5 mg daily during chronic treatment.

Allergic and skin disorders Initial doses of 5-15 mg daily are commonly adequate.

**Collagenosis** Initial doses of 20-30 mg daily are frequently effective. Those with more severe symptoms may require higher doses.

**Rheumatoid arthritis** The usual initial dose is 10-15 mg daily. The lowest daily maintenance dose compatible with tolerable symptomatic relief is recommended.

**Blood disorders and lymphoma** An initial daily dose of 15-60 mg is often necessary with reduction after an adequate clinical or haematological response. Higher doses may be necessary to induce remission in acute leukaemia.

#### **Special populations**

**Use in elderly** Treatment of elderly patients, particularly if long-term, should be planned bearing in mind the more serious consequences of the common side-effects of corticosteroids in old age (see also 'Special warnings and special precautions for use').

#### Use in children

Although appropriate fractions of the actual dose may be used, dosage will usually be determined by clinical response as in adults (see also Section 4.4

'Special warnings and special precautions for use' and Section 4.8 'Undesirable effects'). Alternate day dosage is preferable where possible.

#### Method of administration

Oral Use.

#### 4.3 Contraindications

- Hypersensitivity to prednisolone or any of the excipients (see Section 6.1 'List of Excipients').
- Systemic infections unless specific anti-infective therapy is employed.
- Ocular herpes simplex because of possible perforation.

#### 4.4 Special warnings and precautions for use

Patients/ and or carers should be warned that potentially severe psychiatric adverse reactions may occur with systemic steroids (see Section 4.8 'Undesirable effects'). Symptoms typically emerge within a few days or weeks of starting the treatment. Risks may be higher with high doses/ systemic exposure (see also Section 4.5 'Interaction with other medicinal products and other forms of interaction'), although dose levels do not allow prediction of the onset, type, severity or duration of reactions. Most reactions recover after either dose reduction or withdrawal, although specific treatment may be necessary.

Patients/carers should be encouraged to seek medical advice if worrying psychological symptoms develop, especially if depressed mood or suicidal ideation is suspected. Patients/carers should also be alert to possible psychiatric disturbances that may occur either during or immediately after dose tapering/ withdrawal of systemic steroids, although such reactions have been reported infrequently.

Particular care is required when considering the use of systemic corticosteroids in patients with existing or previous history of severe affective disorders in themselves or in their first degree relatives. These would include depressive or manic depressive illness and previous steroid psychosis.

**Tumorigenicity:** direct tumour-inducing effects of the glucocorticoids are not known, but the particular risk that malignancies in patients undergoing immunosuppression with these or other drugs will spread more rapidly is a well-recognised problem (see Section 4.5 'Interaction with other medicinal products and other forms of interaction').

**Calciphylaxis** may occur very rarely during treatment with corticosteroids (see section 4.8 'Undesirable effects'). Although calciphylaxis is most commonly observed in patients who have end stage kidney failure, it has also been reported in patients taking corticosteroids who have minimal or no renal impairment and normal calcium, phosphate and parathyroid hormone levels. Patients/carers should be advised to seek medical advice if symptoms develop.

Caution is necessary when oral corticosteroids, including Prednisolone, are prescribed in patients with the following conditions, and frequent patient monitoring is necessary:

- Tuberculosis: Those with a previous history of, or X-ray changes characteristic of, tuberculosis. The emergence of active tuberculosis can, however, be prevented by the prophylactic use of anti-tuberculosis therapy.
- Inflammatory bowel disease: Symptoms recurred in a patient with Crohn's disease on changing from conventional to enteric-coated tablets of prednisolone. This was not an isolated occurrence in the author's unit, and it was advocated that only non-enteric coated prednisolone tablets should be used in Crohn's disease, and that the enteric coated form should be used with caution in any condition characterized by diarrhoea or a rapid transit time.
- Hypertension.
- Congestive heart failure.
- Liver failure.
- Hepatic disease: In patients with acute and active hepatitis, protein binding of the glucocorticoids will be reduced and peak concentrations of administered glucocorticoids increased. Elimination of prednisolone will also be impaired. There is an enhanced effect of corticosteroids in patients with cirrhosis.
- Renal insufficiency.
- Diabetes mellitus or in those with a family history of diabetes.
- Osteoporosis: This is of special importance in post-menopausal females who are at particular risk.
- Corticosteroid requirements may be reduced in menopausal and postmenopausal women.
- Patients with a history of severe affective disorders and particularly those with a previous history of steroid-induced psychoses.
- Also, existing emotional instability or psychotic tendencies may be aggravated by corticosteroids including prednisolone.
- Epilepsy, and/or seizure disorders.
- Peptic ulceration.
- Previous steroid myopathy.
- Glucocorticoids should be used cautiously in patients with myasthenia gravis receiving anticholinesterase therapy.

- Because cortisone has been reported rarely to increase blood coagulability and to precipitate intravascular thrombosis, thromboembolism, and thrombophlebitis, corticosteroids should be used with caution in patients with thromboembolic disorders.
- Duchenne's muscular dystrophy: transient rhabdomyolysis and myoglobinuria may occur following strenuous physical activity. It is not known whether this is due to prednisolone itself or the increased physical activity.

Undesirable effects may be minimised by using the lowest effective dose for the minimum period and by administering the daily requirement as a single morning dose on alternate days. Frequent patient review is required to titrate the dose appropriately against disease activity (see Section 4.2 'Posology and method of administration').

Adrenocortical Insufficiency: Pharmacologic doses of corticosteroids administered for prolonged periods may result in hypothalamic-pituitaryadrenal (HPA) suppression (secondary adrenocortical insufficiency). The degree and duration of adrenocortical insufficiency produced is variable among patients and depends on the dose, frequency, time of administration, and duration of glucocorticoid therapy.

In addition, acute adrenal insufficiency leading to a fatal outcome may occur if glucocorticoids are withdrawn abruptly. Drug induced secondary adrenocortical insufficiency may therefore be minimized by gradual reduction of dosage. This type of relative insufficiency may persist for months after discontinuation of therapy; therefore, in any situation of stress occurring during that period, hormone therapy should be reinstituted. Since mineralocorticoid secretion may be impaired, salt and/or a mineralocorticoid should be administered concurrently. During prolonged therapy any intercurrent illness, trauma, or surgical procedure will require a temporary increase in dosage; if corticosteroids have been stopped following prolonged therapy they may need to be temporarily re-introduced.

Patients should carry "Steroid treatment" cards which give clear guidance on the precautions to be taken to minimise risk and which provide details of prescriber, drug, dosage and the duration of treatment.

Anti-inflammatory/Immunosuppressive effects and Infection: Suppression of the inflammatory response and immune function increases the susceptibility to infections and their severity. The clinical presentation may often be atypical and serious infection such as septicaemia and tuberculosis may be masked and may reach an advanced stage before being recognised when corticosteroids including prednisolone are used. The immunosuppressive effects of glucocorticoids may result in activation of latent infection or exacerbation of intercurrent infections.

**Chickenpox:** Chickenpox is of particular concern since this normally minor illness may be fatal in immunosuppressed patients. Patients (or parents of children) without a definite history of chickenpox should be advised to avoid close personal contact with chickenpox or herpes zoster and if exposed they should seek urgent medical attention. Passive immunisation

with varicella-zoster immunoglobulin (VZIG) is needed by exposed nonimmune patients who are receiving systemic corticosteroids or who have used them within the previous 3 months; this should be given within 10 days of exposure to chickenpox. If a diagnosis of chickenpox is confirmed, the illness warrants specialist care and urgent treatment.

Corticosteroids should not be stopped and the dose may need to be increased.

**Measles:** Patients should be advised to take particular care to avoid exposure to measles, and to seek immediate medical advice if exposure occurs. Prophylaxis with intramuscular normal immunoglobulin may be needed.

Administration of Live Vaccines: Live vaccines should not be given to individuals on high doses of corticosteroids, due to impaired immune response. Live vaccines should be postponed until at least 3 months after stopping corticosteroid therapy.

(See also Section 4.5 'Interaction with other medicinal products and other forms on interaction').

**Ocular Effects:** Prolonged use of corticosteroids may produce posterior subcapsular cataracts and nuclear cataracts (particularly in children), exophthalmos, or increased intraocular pressure, which may result in glaucoma with possible damage to the optic nerves. Establishment of secondary fungal and viral infections of the eye may also be enhanced in patients receiving glucocorticoids.

Corticosteroids should be used cautiously in patients with ocular herpes simplex because of possible perforation.

Systemic glucocorticoid treatment can cause severe exacerbation of bullous exudative retinal detachment and lasting visual loss in some patients with idiopathic central serous chorioretinopathy (See Section 4.8 'Undesirable effects').

**Visual disturbance:** Visual disturbance may be reported with systemic and topical corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes which may

include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids.

**Cushing's disease:** Because glucocorticoids can produce or aggravate Cushing's syndrome, glucocorticoids should be avoided in patients with Cushing's disease

There is an enhanced effect of corticosteroids in patients with hypothyroidism. Psychic derangements may appear when corticosteroids, including prednisolone, are used, ranging from euphoria, insomnia, mood swings, personality changes, and severe depression, to frank psychotic manifestations (see Section 4.8 'Undesirable effects').

**Raised intracranial pressure:** Raised intracranial pressure with papilloedema (pseudotumour cerebri) associated with corticosteroid treatment has been reported in both children and adults. The onset usually occurs after treatment withdrawal (See section 4.8 'Undesirable effects').

**Scleroderma renal crisis:** Caution is required in patients with systemic sclerosis because of an increased incidence of (possibly fatal) scleroderma renal crisis with hypertension and decreased urinary output observed with a daily dose of 15 mg or more prednisolone. Blood pressure and renal function (s-creatinine) should therefore be routinely checked. When renal crisis is suspected, blood pressure should be carefully controlled.

#### Use in the elderly

Treatment of elderly patients, particularly if long term, should be planned bearing in mind the more serious consequences of the common side-effects of corticosteroids in old age, especially osteoporosis, diabetes, hypertension, hypokalaemia, susceptibility to infection and thinning of the skin. Close clinical supervision is required to avoid life threatening reactions.

#### **Paediatric population**

Corticosteroids cause growth retardation in infancy, childhood and adolescence, which may be irreversible, and therefore long-term administration of pharmacological doses should be avoided. If prolonged therapy is necessary, treatment should be limited to the minimum suppression of the hypothalamo-pituitary adrenal axis and growth retardation. The growth and development of infants and children should be closely monitored. Treatment should be administered where possible as a single dose on alternate days.

There is an increased risk of nuclear cataracts (see Section 4.8 'Undesirable Effects).

#### Important information regarding the ingredients

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Sodium: This medicinal product contains less than 1 mmol of sodium (23mg) per tablet that is to say essentially 'sodium free'.

#### 4.5 Interaction with other medicinal products and other forms of interaction

Vaccines	Live vaccines should not be given to individuals with impaired immune responsiveness. The antibody response to other vaccines may be diminished.		
Antacids	The absorption of prednisolone may be reduced by large doses of some antacids such as magnesium trisilicate or aluminium hydroxide.		
Antibacterials	Rifamycin accelerate metabolism of corticosteroids and thus may reduce their effect. Erythromycin inhibits metabolism of methylprednisolone and possibly other corticosteroids. Prednisolone can lower plasma levels of isoniazid. Where a reduced response during concurrent use is noted, dosage adjustment of isoniazid may be necessary.		
Anticoagulants	Response to anticoagulants may be reduced or, less often, enhanced by corticosteroids. Close monitoring of the INR or prothrombin time is required to avoid spontaneous bleeding.		
Antidiabetic agents	Glucocorticoids may increase blood glucose levels. Patients with diabetes mellitus receiving concurrent insulin and/or oral hypoglycemic agents may require dosage adjustments of such therapy.		
Antiepileptics	Carbamazepine, phenobarbital, phenytoin, and primidone accelerate metabolism of corticosteroids and may reduce their effect.		
Antifungals	Risk of hypokalaemia may be increased with amphotericin, therefore concomitant use with corticosteroids should be avoided unless corticosteroids are required to control reactions; ketoconazole inhibits metabolism of		

	methylprednisolone and possibly other corticosteroids.		
Antimuscarinics (Anticholinergics)	Prednisolone has been shown to have antimuscarinic activity. If used in combination with another antimuscarinic drug could cause impairment to memory and attention in the elderly.		
Antithyroids	Prednisolone clearance increased by the use of carbimazole and thiamazole.		
Cardiac Glycosides	Increased toxicity if hypokalaemia occurs with corticosteroids.		
Ciclosporin	Concomitant administration of prednisolone and ciclosporin may result in decreased plasma clearance of prednisolone (i.e. increased plasma concentration of prednisolone). The need for appropriate dosage adjustment should be considered when these drugs are administered concomitantly.		
Cytotoxics	Increased risk of haematological toxicity with methotrexate.		
Hepatic microsomal enzyme inducers	Drugs that induce hepatic enzyme cytochrome P-450 (CYP) isoenzyme 3A4 such as phenobarbital, phenytoin, rifampicin, rifabutin, carbamazepine, primidone and aminoglutethimide may reduce the therapeutic efficacy of corticosteroids by increasing the rate of metabolism. Lack of expected response may be observed and dosage of Prednisolone may need to be increased.		
Hepatic microsomal enzyme inhibitors	Co-treatment with CYP3A inhibitors, including cobicistat-containing products, is expected to increase the risk of systemic side-effects. The combination should be avoided unless the benefit outweighs the increased risk of systemic corticosteroid side-effects, in which case patients should be monitored for systemic corticosteroid side-effects. Drugs that inhibit hepatic enzyme cytochrome P-450 (CYP) isoenzyme 3A4 (e.g. ketoconazole, troleandomycin) may decrease glucocorticoid clearance. Dosages of glucocorticoids given in combination with such drugs may need to be decreased to avoid potential adverse effects		
Hormonal	Oral contraceptives increased prednisolone concentrations by 131%.		

contraceptives	May increase AUC and reduce clearance in oral				
	contraceptives containing ethinylestradiol, mestranol,				
	desogestrel, levonorgestrel, norgestrel or				
	norethisterone.				
	Tumorigenicity: direct tumour-inducing effects of the				
	glucocorticoids are not known, but the particular risk				
<b>Immunosuppressants</b>	that malignancies in patients undergoing				
	immunosuppression with these or other drugs will				
	spread more rapidly is a well-recognised problem.				
Liquorice	Glycyrrhizin can delay the clearance of prednisolone				
Mifepristone	Effect of corticosteroids may be reduced for 3-4 days				
	after mifepristone				
	Concomitant administration of ulcerogenic drugs such				
	as indomethacin during corticosteroid therapy may				
	increase the risk of GI ulceration. Aspirin should be				
	used cautiously in conjunction with glucocorticoids in				
	patients with hypoprothrombinaemia. Although				
	concomitant therapy with salicylate and				
	corticosteroids does not appear to increase the				
	incidence or severity of GI ulceration, the possibility				
Non-steroidal anti-	of this effect should be considered.				
inflammatory drugs	Serum salicylate concentrations may decrease when				
	corticosteroids are administered concomitantly. The				
	renal clearance of salicylates is increased by				
	corticosteroids and steroid withdrawal may result in				
	salicylate intoxication. Salicylates and corticosteroids				
	should be used concurrently with caution. Patients				
	receiving both drugs should be observed closely for				
	adverse effects of either drug.				
	Oestrogens may potentiate the effects of				
Oestrogens	glucocorticolds and dosage adjustments may be				
5	required it oestrogens are added to or withdrawn from				
	a stable dosage regimen.				
	Ritonavir possibly increases plasma concentrations of				
Duotoggo inhibitoug	prednisolone and other corticosteroids by reduction in				
r roleuse innibilors	clearance of prednisolone through the inhibition of				
	P450 isoenzyme CYP3A4.				
	The desired effects of hypoglycaemic agents				
Othor	(including insulin), antihypertensives and diuretics are				
	antagonised by corticosteroids; and the hypokalaemic				
	effect of acetazolamide, loop diuretics, thiazide				

	diuretics, carbenoxolone and theophylline are enhanced.		
Somatropin	Growth promoting effect may be inhibited.		
Sympathomimetics	Increased risk of hypokalaemia if high doses of corticosteroids given with high doses of bambuterol, fenoterol, formoterol, ritodrine, salbutamol, salmeterol and terbutaline.		

#### 4.6 Fertility, pregnancy and lactation

#### Pregnancy

The ability of corticosteroids to cross the placenta varies between individual drugs, however, 88% of prednisolone is inactivated as it crosses the placenta. Administration of corticosteroids to pregnant animals can cause abnormalities of foetal development including cleft palate, intra-uterine growth retardation and effects on brain growth and development. There is no evidence that corticosteroids result in an increased incidence of congenital abnormalities, such as cleft palate/lip in man. However, when administered for prolonged periods or repeatedly during pregnancy, corticosteroids may increase the risk of intra-uterine growth retardation. Hypoadrenalism may, in theory, occur in the neonate following prenatal exposure to corticosteroids but usually resolves spontaneously following birth and is rarely clinically important. Cataracts have been observed in infants born to mothers treated with long-term prednisolone during pregnancy. As with all drugs, corticosteroids should only be prescribed when the benefits to the mother and child outweigh the risks. When corticosteroids are essential however, patients with normal pregnancies may be treated as though they were in the non-gravid state.

Patients with pre-eclampsia or fluid retention require close monitoring.

#### **Breast-feeding**

Corticosteroids are excreted in small amounts in breast milk. Corticosteroids distributed into breast milk may suppress growth and interfere with endogenous glucocorticoid production in nursing infants. Since adequate reproductive studies have not been performed in humans with glucocorticoids, these drugs should be administered to nursing mothers only if the benefits of therapy are judged to outweigh the potential risks to the infant.

The concentration of the steroid in the milk can be between 5 and 25% of those in the serum and the two roughly parallel one another after an oral dose.

There are no reports found regarding neonatal toxicity following exposure to corticosteroids during lactation, however if maternal doses >40mg/day of prednisolone is prescribed, the infant should be monitored for adrenal suppression.

#### 4.7 Effects on ability to drive and use machines

The effect of Prednisolone on the ability to drive or use machinery has not been evaluated. There is no evidence to suggest that prednisolone may affect these abilities.

#### 4.8 Undesirable effects

A wide range of psychiatric reactions including affective disorders (such as irritable, euphoric, depressed and labile mood, and suicidal thoughts), psychotic reactions (including mania, delusions, hallucinations, and aggravation of schizophrenia), behavioural disturbances, irritability, anxiety, sleep disturbances, and cognitive dysfunction including confusion and amnesia have been reported. Reactions are common and may occur in both adults and children. In adults, the frequency of severe reactions has been estimated to be 5-6%. Psychological effects have been reported on withdrawal of corticosteroids; the frequency is unknown.

The incidence of predictable undesirable effects, including hypothalamicpituitary-adrenal suppression correlates with the relative potency of the drug, dosage, timing of administration and the duration of treatment (see special warnings for use).

Undesirable effects are listed by MedDRA System Organ Classes. Assessment of undesirable effects is based on the following frequency groupings:

- Very common:  $\geq 1/10$
- Common:  $\geq 1/100$  to < 1/10
- Uncommon:  $\geq 1/1,000$  to < 1/100
- Rare:  $\geq 1/10,000$  to < 1/1,000
- Very rare: <1/10,000
- Not known: cannot be estimated from the available data

#### Tabulated list of adverse reactions:

System organ class	Frequency	Adverse reaction
Infections and	Not known	Increases susceptibility to, and severity of
Infestations		infections <sup>1</sup> , opportunistic infections,

		recurrence of dormant tuberculosis <sup>2</sup> ,
		oesophageal candidiasis.
Blood and		
lymphatic system	Not known	Leucocytosis.
disorders		
Immune system	Not known	Hypersensitivity including anaphylaxis,
disorders		Scleroderma renal crisis <sup>10</sup> .
	Not known	Suppression of the hypothalamo-pituitary
Endocrine		adrenal axis <sup>3</sup> , cushingoid facies, impaired
disorders		carbohydrate tolerance with increased
		requirement for antidiabetic therapy,
		manifestation of latent diabetes mellitus.
		Sodium and water retention, hypokalaemic
		alkalosis, potassium loss, negative nitrogen
		and calcium balance, glucose intolerance
Metabolism and	Not known	and protein catabolism. Increase both high
nutrition		and low density lipoprotein cholesterol
disorders		concentration in the blood. Increased
		appetite <sup>4</sup> . Weight gain, obesity,
		hyperglycaemia, dyslipidaemia.
	Very rare	
		Irritability, depressed and labile mood,
		suicidal thoughts, psychotic reactions,
	C	mania, delusions, hallucinations, and
Psychiatric	Common	aggravation of schizophrenia. behavioural
disorders		disturbances, anxiety, sleep disturbances,
		and cognitive dysfunction including
		Confusion and amnesia.
	Not known	Euphoria, psychological dependence,
		Depression incompio digginogg headache
	Not known	Depression, insomina, dizziness, neadache,
Nervous system		papilloedema (nseudotumor cerebri) <sup>6</sup>
disorders		Aggregation of englancy, endural
		linomatosis, vertebrohasilar stroke <sup>7</sup>
		Glaucoma papilloedema posterior
Eye disorders	Not known	subcansular cataracts, nuclear cataracts
		(particularly in children) exophthalmos
		corneal or scleral thinning, exacerbation of
		onhthalmic viral or fungal disease
		Severe exacerbation of bullous evudative
		ratinal detachment: lasting visual loss in
		some notionts with idionsthic control account
		some patients with idiopathic central serous

		chorioretinopathy.8
	Not known	Vision, blurred (see section 4.4)
Ear and labyrinth disorders	Not known	Vertigo.
Cardiac disorders	Not known	Congestive heart failure in susceptible patients, hypertension, increased risk of heart failure. Increased risk of cardiovascular disease, including myocardial infarction <sup>9</sup> , Bradycardia*
Vascular disorders	Not known	Thromboembolism.
Gastrointestinal disorders	Not known	Dyspepsia, nausea, peptic ulceration with perforation and haemorrhage, abdominal distension, abdominal pain, diarrhoea, oesophageal ulceration, acute pancreatitis.
Skin and subcutaneous tissue disorders	Not known	Hirsutism, skin atrophy, bruising, striae, telangiectasia, acne, increased sweating, pruritis, rash, urticaria.
Musculoskeletal and connective tissue disorders	Not known	Proximal myopathy, osteoporosis, vertebral and long bone fractures, avascular osteonecrosis, tendon rupture, tendinopathies (particularly of the Achilles and patellar tendons), myalgia, growth suppression in infancy, childhood and adolescence.
Reproductive system and breast disorders	Not known	Menstrual irregularity, amenorrhoea.
General disorders and administration site conditions	Not known	Fatigue, malaise, impaired healing
Investigations	Not known	Increased intra-ocular pressure, may suppress reactions to skin tests.

1. With suppression of clinical symptoms and signs.

2. See Section 4.4 'Special warnings and precautions for use'.

3. Particularly in times of stress, as in trauma, surgery or illness.

4. Which may result in weight gain.

5. See Section 4.4 'Special warnings and precautions for use'.

6. Usually after treatment withdrawal.

7. Exacerbation of giant cell arteritis, with clinical signs of evolving stroke has been attributed

to prednisolone.

8. See Section 4.4 'Special warnings and precautions for use'.

9. With high dose therapy.

10. Scleroderma renal crisis: Amongst the different subpopulations the occurrence of scleroderma renal crisis varies. The highest risk has been reported in patients with diffuse systemic sclerosis. The lowest risk has been reported in patients with limited systemic sclerosis (2%) and juvenile onset systemic sclerosis (1%) \*Following high dose

#### Withdrawal symptoms and signs

Too rapid a reduction of corticosteroid dosage following prolonged treatment can lead to acute adrenal insufficiency, hypotension and death (see 'Special warnings and precautions for use' and Section 4.2 'Posology and method of administration'). A steroid "withdrawal syndrome" seemingly unrelated to adrenocortical insufficiency may also occur following abrupt discontinuance of glucocorticoids. This syndrome includes symptoms such as: anorexia, nausea, vomiting, lethargy, headache, fever, joint pain, desquamation, myalgia, arthralgia, rhinitis, conjunctivitis, painful itchy skin nodules, weight loss, and/or hypotension. These effects are thought to be due to the sudden change in glucocorticoid concentration rather than to low corticosteroid levels. Psychological effects have been reported on withdrawal of corticosteroids.

#### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme Website: <u>www.mhra.gov.uk/yellowcard</u> or search for MHRA Yellow Card in the Google Play or Apple App Store.

#### 4.9 Overdose

Reports of acute toxicity and/or death following overdosage of glucocorticoids are rare. In the event of overdosage no specific antidote is available. Treatment is supportive and symptomatic. Serum electrolytes should be monitored.

High systemic doses of corticosteroids caused by chronic use have been associated with adverse effects such as neuropsychiatric disorders (psychosis, depression, hallucinations), cardiac dysrhythmias and Cushing's syndrome.

# 5 PHARMACOLOGICAL PROPERTIES

#### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Corticosteroids for systemic use.

ATC code: H02AB06

Naturally occurring glucocorticoids (hydrocortisone and cortisone), which also have salt- retaining properties, are used as replacement therapy in adrenocortical deficiency states. Their synthetic analogs are primarily used for their potent anti-inflammatory effects in disorders of many organ systems.

Glucocorticoids cause profound and varied metabolic effects. In addition, they modify the body's immune responses to diverse stimuli.

#### 5.2 Pharmacokinetic properties

Prednisolone is rapidly and apparently almost completely absorbed after oral administration; it reaches peak plasma concentrations after 1-3 hours. There is however wide inter-subject variation suggesting impaired absorption in some individuals. Plasma half-life is about 3 hours in adults and somewhat less in children. Its initial absorption, but not its overall bioavailability, is affected by food. Prednisolone has a biological half-life lasting several hours, making it suitable for alternate-day administration regimens.

Although peak plasma prednisolone levels are somewhat lower after administration of Prednisolone and absorption is delayed, total absorption and bioavailability are the same as after plain prednisolone. Prednisolone shows dose dependent pharmacokinetics, with an increase in dose leading to an increase in volume of distribution and plasma clearance. The degree of plasma protein binding determines the distribution and clearance of free, pharmacologically active drug. Reduced doses are necessary in patients with hypoalbuminaemia.

Prednisolone is metabolised primarily in the liver to a biologically inactive compound. Liver disease prolongs the half-life of prednisolone and, if the patient has hypoalbuminaemia, also increases the proportion of unbound drug and may thereby increase adverse effects.

Prednisolone is excreted in the urine as free and conjugated metabolites, together with small amounts of unchanged prednisolone.

Significant differences in the pharmacokinetics of prednisolone amongst menopausal women have been described. The postmenopausal women had reduced unbound clearance (30%), reduced total clearance and increased half-life of prednisolone.

#### 5.3 Preclinical safety data

Not applicable.

# 6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients Lactose Maize Starch Colloidal Silicon Dioxide Sodium Starch Glycollate Stearic Acid Magnesium Stearate.

# 6.2 Incompatibilities None

#### 6.3 Shelf life

Container : 36 months. Blister : 24 months

#### 6.4 Special precautions for storage

Container : Do not store above 25°C. Store in the original container. Keep the container tightly closed.

Blister packs: Do not store above 25°C. Store in the original package.

#### 6.5 Nature and contents of container

Polypropylene tubes fitted with polyethylene cap. Pack sizes: 100, 250, 500, 1000 and 5000 tablets.

Blister of PVC / blister aluminium foil Pack size: 28 tablets

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling Not applicable

# 7 MARKETING AUTHORISATION HOLDER

Bristol Laboratories Ltd Unit 3, Canalside, Northbridge Road, Berkhamsted, Hertfordshire HP4 1EG United Kingdom

# 8 MARKETING AUTHORISATION NUMBER(S) PL 17907/0455 PL 17907/0456

# 9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first Authorisation: 1<sup>st</sup> March 2005 Date of Renewal of Authorisation: 13<sup>th</sup> May 2008

# **10 DATE OF REVISION OF THE TEXT**

18/07/2022